
Figure 1A


Effect of different supernatants of lactic acid bacteria on the formation of capillary structures by HUVEC cells



- 1. Supernatant of lactic acid bacteria
- Concentrated supernatant of lactic acid bacteria (molecules > 5 000 kDa)
- 3. Filtrate of lactic acid bacteria (molecules ≤ à 5 000 kDa)

Figure 1B


Effect of different supernatants of lactic acid bacteria on the formation of capillary structure by HUVEC cells



- 1. Supernatant of lactic acid bacteria
- 2. Concentrated supernatant of lactic acid bacteria (molecules > 5 000 kDa)
- 3. Filtrate of lactic acid bacteria (molecules ≤ à 5 000 kDa)

Figure 2A

Effect of different BIO-K PLUS supernatants on the formation of the capillary structure by HUVEC cells




- Supernatant of BIO-K PLUS (with milk product) 1.
- Supernatant of BIO-K PLUS (without milk product) Supernatant of BIO-K PLUS (with milk product) pH7.2 2.
- Supernatant of BIO-K PLUS (without milk product) pH7.2

4 / 36

Figure 2B


Effect of different BIO-K PLUS supernatants on the formation of the capillary structure by HUVEC cells



- 1. Supernatant of BIO-K PLUS (with milk product)
- 2. Supernatant of BIO-K PLUS (without milk product)
- Supernatant of BIO-K PLUS (with milk product) pH7.2
 Supernatant of BIO-K PLUS (without milk product) pH7.2

Figure 3A


Effect of lactic acid bacteria supernatant on the migration of BAEC cells



- 1. Supernatant of lactic acid bacteria
- 2. Concentrated supernatant of lactic acid bacteria (molecules > 5 000 kDa)
- 3. Filtrate of lactic acid bacteria (molecules ≤ à 5 000 kDa)

Figure 3B


Effect of BIO-K PLUS supernatant on the migration of BAEC cells



- BIO-K PLUS supernatant (with milk product)
 BIO-K PLUS supernatant (without milk product)

Figure 4A


Effect of lactic acid bacteria supernatant on migration of HUVEC cells



- 1. Supernatant of lactic acid bacteria
- 2. Concentrated supernatant of lactic acid bacteria (molecules > 5 000 kDa)
- 3. Filtrate of lactic acid bacteria (molecules ≤ à 5 000 kDa)

Figure 4B

Effect of BIO-K PLUS supernatant on migration of HUVEC cells



- 1. Supernatant of BIO-K PLUS (with milk product)
- 2. Supernatant of BIO-K PLUS (without milk product)

Figure 5A

Effect of BIO-K PLUS supernatant (with milk product) on migration of BAEC cells




Figure 5B

Effect of BIO-K PLUS supernatant (without milk product) on migration of BAEC cells




Figure 6A

Effect of BIO-K PLUS supernatant (with milk product) on HUVEC cells proliferation





Figure 6B


Effect of BIO-K PLUS supernatant (without milk product) on HUVEC cells proliferation



Incubation time

Figure 7A


Effect of BIO-K PLUS supernatant (with milk product) on the proliferation of HUVEC cells induced by bFGF



Incubation time

Figure 7B

Effect of BIO-K PLUS supernatant (without milk product) on the proliferation of HUVEC cells induced by bFGF



PCT/CA2004/001968 WO 2005/046703

15 / 36

Figure 8A

Effect of lactic bacteria supernatants on the proliferation of endothelial cells after 65 h treatment (n =1)





Figure 8B









Figure 9



-VEGF



SUBSTITUTE SHEET (RULE 26)



* Figure 11A






Figure 11B








PCT/CA2004/001968 WO 2005/046703


22 / 36

Figure 12












FIGURE 15





Figure 17

Effect of different supernatants of BIOK PLUS on the formation of capillary structures by endothelial cells (HUVEC)






FIGURE 18

inhibition of endothelial cells Effect of Bio-K Plus supernatants on cel




FIGURE 1

Effect of Bio-K Plus supernatants on BAECs cell migration






FIGURE 20




Effect of *Bio-K Plus* supernatant on VEGFR-2 phosphorylation



IP: VEGFR-2 ID: p(Tyr)

Ctl: Control
BKL: BK milk
BK: BK without milk

phosphorylation of protéine Erk by Effect of Bio-K Plus supernatant VEGF



Ctl: Control BK: BK without milk

FIGURE 23

Antagonist effect of Bio-K Plus supernatants on different receptors of growth factors





FIGURE 2

High throughput screening of growth factor



Molecular screening of receptors antagonists

